Author Index

Afroze, T., see Shippen-Lentz, D., 113
Anders, R.F., see Foley, M., 69
Anders, R.F., see Lustigman, S., 261
Armah, G.E., Nishikawa, S., Miki, S., Omata, Y., Nakabayashi, T. and Tomita, K.-i.
Conformation and immunogenicity of engineered repeating segment of the circumsporozoite surface protein of *Plasmodium falciparum*, 135
Ashall, F.
Characterisation of an alkaline peptidase of *Trypanosoma cruzi* and other trypanosomatids, 77
Ayala, P., see Samuelson, J., 281

Bacic, A., see McConville, M.J., 57
Barker, G.C., see Mercer, J.G., 89
Beach, D.H., Holz, Jr., G.G., Singh, B.N. and Lindmark, D.G.
Fatty acid and sterol metabolism of cultured *Trichomonas vaginalis* and *Tritrichomonas foetus*, 175
Bishop, R.P.
Extensive homologies between *Leishmania donovani* chromosomes of markedly different size, 1
Bouhours, J.-F., see Persat, F., 97
Brown, G.V., see Lustigman, S., 261
Button, L.L., see Frommel, T.O., 25

Cazzulo, J.J., Hellman, U., Couso, R. and Parodi, A.J.A.
Amino acid and carbohydrate composition of a lysosomal cysteine proteinase from *Trypanosoma cruzi*. Absence of phosphorylated mannose residues, 41
Cochrane, A.H., see Kilejian, A., 291
Coggins, L., see Hall, R., 253
Comley, J.C.W., see Hayes, D.J., 159
Coppel, R.L., see Lustigman, S., 261
Cornelissen, A.W.C.A., see Köck, J., 295
Couso, R., see Cazzulo, J.J., 41

De Boeck, H., see Zidovetzki, R., 33
Deguercy, A., Hommel, M. and Schrével, J.
Purification and characterization of 37-kilodalton proteases from *Plasmodium falciparum* and *Plasmodium berghei* which cleave erythrocyte cytoskeletal components, 233
Dunne, D.W., see Havercroft, J.C., 211

Foley, M., Murray, L.J. and Anders, R.F.
The ring-infected erythrocyte surface antigen protein of *Plasmodium falciparum* is phosphorylated upon association with the host cell membrane, 69
Frommel, T.O., Button, L.L., Fujikura, Y. and McMaster, W.R.
The major surface glycoprotein (GP63) is present in both life stages of *Leishmania*, 25
Fujikura, Y., see Frommel, T.O., 25

Gould, H., see Kara, U., 19

Hall, R., Coggins, L., McKellar, S., Shiels, B. and Tait, A.
Characterisation of an extrachromosomal DNA element from *Theileria annulata*, 253
Harris, B.G., see Srinivasan, N.G., 149
Havercroft, J.C., Huggins, M.C., Dunne, D.W. and Taylor, D.W.
Characterisation of Sm20, a 20-kilodalton calcium-binding protein of *Schistosoma mansoni*, 211
Hayes, D.J., O'Dowd, A.B., Selwood, D.L., Stables, J.N., Comley, J.C.W. and Rees, M.J.
The effect of two novel analogues of antimycin A on oxygen consumption and survival of filarial nematodes in vitro, 159
Hellman, U., see Cazzulo, J.J., 41
Hommel, M., see Deguercy, A., 233
Holz, Jr., G.G., see Beach, D.H., 175
Howells, R.E., see Mercer, J.G., 89
Huggins, M.C., see Havercroft, J.C., 211

Jagannatha Rao, G.S., see Srinivasan, N.G., 149

Kara, U., Murray, B., Pam, C., Lahnstein, J., Gould, H., Kidson, C. and Saul, A.
Chemical characterization of the parasitophorous vacuole membrane antigen QF116 from *Plasmodium falciparum*, 19
Kemp, D.J., see Shirley, M.W., 169
Kidson, C., see Kara, U., 19
Kilejian, A., Yang, Y.-F., Cochrane, A.H. and Rashid, M.A.
Homologous sequences in Plasmodium cynomolgi and the gene of the histidine-rich knob protein of *Plasmodium falciparum (Short Communication)*, 291
Kilejian, A., see Rashid, M.A., 49
Köck, J. and Cornelissen, A.W.C.A.
The 5S ribosomal RNA genes of *Crithidia fasciculata (Short Communication)*, 295

Lahnstein, J., see Kara, U., 19
Lima, M.F. and Villalta, F.
Trypanosoma cruzi receptors for human transferrin and their role, 245
Limberger, R.J. and McReynolds, L.A.
Filarial paramyosin: cDNA sequences from *Dirofilaria immitis* and *Onchocerca volvulus*, 271
Lindmark, D.G., see Beach, D.H., 175
Lubega, G.W. and Prichard, R.K.
Specific interaction of benzimidazole anthelmintics with tubulin: high-affinity binding and benzimidazole resistance in *Haemonchus contortus*, 221
Lustigman, S., Anders, R.F., Brown, G.V. and Coppel, R.L.
The mature-parasite-infected erythrocyte surface antigen (MESA) of *Plasmodium falciparum* associated with the erythrocyte membrane skeletal protein, band 4.1, 261

Maguire, P.A. and Sherman, I.W.
Phosopholipid composition, cholesterol content and choles-

terol exhange in *Plasmodium falciparum*-infected red cells, 105
Maguire, P.A., see Zidovetzki, R., 33
McConville, M.J. and Bacic, A.
The glycoinositolphospholipid profiles of two *Leishmania major* strains that differ in lipophosphoglycan expression, 57
McKellar, S., see Hall, R., 253
McMaster, W.R., see Frommel, T.O., 25
McReynolds, L.A., see Limberger, R.J., 271
Mercer, J.G., Barker, G.C., Howells, R.E. and Rees, H.H.
Investigation of ecdysteroid excretion by adult *Dirofilaria immitis* and *Brugia pahangi*, 89
Miki, S., see Armah, G.E., 135
Mojon, M., see Persat, F., 97
Murray, B., see Kara, U., 19
Murray, L.J., see Foley, M., 69
Nakabayashi, T., see Armah, G.E., 135

Nishikawa, S., see Armah, G.E., 135

O'Dowd, A.B., see Hayes, D.J., 159
Omata, Y., see Armah, G.E., 135
Orozco, E., see Samuelson, J., 281

Pallister, J., see Shirley, M.W., 169
Pam, C., see Kara, U., 19
Parodi, A.J.A., see Cazzulo, J.J., 41
Patton, C.L., see Wiser, M.F., 121
Pereira, M.E.A., see Prioli, R.P., 191
Persat, F., Bouhours, J.-F., Mojon, M. and Petavy, A.-F.
Glycosphingolipids of *Echinococcus multilocularis* metacestodes, 97
Petavy, A.-F., see Persat, F., 97
Prichard, R.K., see Lubega, G.W., 221
Prioli, R.P., Rosenberg, I. and Pereira, M.E.A.
High- and low-density lipoproteins enhance infection of *Trypanosoma cruzi* in vitro, 191
Prowse, S.J., see Shirley, M.W., 169

Rashid, M.A., Yang, Y.-F. and Kilejian, A.
Expression, partial purification and immunogenicity of fragments of the knob protein of *Plasmodium falciparum*, 49
Rashid, M.A., see Kilejian, A., 291
Rees, H.H., see Mercer, J.G., 89
Rees, M.J., see Hayes, D.J., 159
Rosenberg, I., see Prioli, R.P., 191

Samuelson, J., Ayala, P., Orozco, E. and Wirth, D.
Emetine-resistant mutants of *Entamoeba histolytica* overexpress mRNAs for multidrug resistance, 281
Sartorelli, A.C., see Wiser, M.F., 121
Saul, A., see Kara, U., 19
Sauma, S.Y. and Strand, M.
Identification and characterization of glycosylphosphatidylinositol-linked *Schistosoma mansoni* adult worm immunogens, 199
Schrével, J., see Deguercy, A., 233
Selwood, D.L., see Hayes, D.J., 159
Sherman, I.W., see Maguire, P.A., 105
Sherman, I.W., see Zidovetzki, R., 33
Shiels, B., see Hall, R., 253
Shippen-Lentz, D., Afroze, T. and Vezza, A.C.
Heterogeneity and expression of the *Plasmodium falciparum* 5.9S ribosomal RNA genes, 113
Shirley, M.W., Kemp, D.J., Pallister, J. and Prowse, S.J.
A molecular karyotype of *Eimeria tenella* as revealed by contour-clamped homogeneous electric field gel electrophoresis, 169
Singh, B.N., see D.H. Beach, D.H., 175
Srinivasan, N.G., Jagannatha Rao, G.S. and Harris, B.G.
Phosphofructokinase from *Dirofilaria immitis*: effect of fructose 2,6-bisphosphate and AMP on the non-phosphorylated and phosphorylated forms of the enzyme, 149
Stables, J.N., see Hayes, D.J., 159
Strand, M., see Sauma, S.Y., 199
Strauss, P.R. and Wang, J.C.
The *TOP2* gene of *Trypanosoma brucei*: a single-copy gene that shares extensive homology with other TOP2 genes encoding eukaryotic DNA topoisomerase II, 141

Tait, A., see Hall, R., 253
Taylor, D.W., see Havercroft, J.C., 211
Tomita, K.-i., see Armah, G.E., 135

Vezza, A.C., see Shippen-Lentz, D., 113
Villalta, F., see Lima, M.F., 245

Walter, R.D., see Wittich, R.-M., 13
Wang, J.C., see Strauss, P.R., 141
Wirth, D., see Samuelson, J., 281
Wiser, M.F., Sartorelli, A.C. and Patton, C.L.
Association of *Plasmodium berghei* proteins with the host erythrocyte membrane: binding to inside-out vesicles, 121
Wittich, R.-M. and Walter, R.D.
Putrescine *N*-acetyltransferase in *Onchocerca volvulus* and *Ascaris suum*, an enzyme which is involved in polyamine degradation and release of *N*-acetylputrescine, 13

Yang, Y.-F., see Kilejian, A., 291
Yang, Y.-F., see Rashid, M.A., 49

Zidovetszki, R., Sherman, I.W., Maguire, P.A. and De Boeck, H.
A nuclear magnetic resonance study of the interactions of the antimalarials chloroquine, quinacrine, quinine and mefloquine with lipids extracted from normal human erythrocytes, 33

Subject Index

Acanthocheilonema viteae, 159
Affinity purification, 199
Amastigote and promastigote expression, 25
Amastigote receptors for human transferrin, 245
Amino acid composition, 41
Amoeba, 281
AMP, 149
Amphiphilic drug, 33
Anthelmintic, 221
Antifilarial, 159
Ascaris suum, 13

Band 4.1, 261
Benzimidazole, 221
Berenil, 13
Blood-stage parasite, 19
Brugia pahangi, 159

Calcium, 211
Calmodulin, 211
Cestode, 95
Cholesterol, 103, 175
Cholesteryl ester, 175
Chromosome polymorphism, 1
Conformation, 133
Crithidia fasciculata, 295
Cysteine peptidase, 75
Cysteine proteinase, 41

Diagnosis of infection, 87
Dirofilaria immitis, 149, 271
DNA, 281
DNA sequence, 139

Ecdyseroid, excretion, 87
Echinococcus multilocularis, 95
Eimeria tenella, 169
Emetine, 281
Endocytosis, receptor-mediated, 245
Energy generation, 159
Entamoeba histolytica, 281
Enzyme kinetics, 75
Epitope mapping, 19
Erythrocyte, 233
Erythrocyte membrane, 119
Erythrocyte membrane skeleton, 261
Extrachromosomal element, 253

Fatty acid, 175
Ferrotransferrin-dependent amastigote growth, 245
Filaria, 87
Fructose, 2,6-bisphosphate, 149

Galactofuranose, 55
Ganglioside, 95
Gene homology, 291
Glycolipid, 55
Glycophosphospingolipid, 175
Glycosphingolipid, 95
Glycosylphosphatidylinositol, 55
Glycosylphosphatidylinositol anchored protein, 199

Heamonchus contortus, 221
HDL, 191
Histamine, 13
Histidine-rich knob protein, 291
Host, 95

Immunogenicity, 133
Immunoprecipitation, 199
Infection, enhancement, 191
Invasion, 67
Iron delivery from ferrotransferrin, 245

Kinetoplastida, 295
Knob protein, 49

LDL, 191
Leishmania, 25
Leishmania donovani, 1
Leishmania major, 55
Lipid membrane, 33
Lipophosphoglycan, 55
Lipoprotein, 191
Lysosomal enzyme, 41

Major surface glycoprotein, 25
Malaria vaccine, 133
Malaria, 103
Malarial antibody, 49
Malarial parasite, 119
Mechanism of benzimidazole resistance, 221
Molecular karyotype, 169
Monoclonal antibody, 199
Multidrug resistance, 281
Mutant, 281
Myristylation, 19

N-acetylation, 13
NMR, 33
Nucleotide, 159
Nucleotide sequence, 271

Oligosaccharide composition, 41
Onchocerca volvulus, 13, 271
Orthogonal field alternation gel electrophoresis, 1
Oxygen consumption, 159

Paramyosin, 271
Parasitophorous vacuole membrane, 19

Peptidase, 75
Phosphatidylinositol-specific phospholipase C, 199
Phosphofructokinase, 149
Phospholipid, 103, 175
Phosphorylation, 67, 149
Physiological assay, 149
Plasmodium berghei, 119, 233
Plasmodium cynomolgi, 291
Plasmodium falciparum, 19, 49, 67, 103, 111, 133, 233, 261, 291
Polyamine, 13
Polymerase chain reaction, 281
Protease, 233
Protein-protein interaction, 261
Pulsed field gel electrophoresis, 169
Putrescine, 13

Recombinant, 271
Recombinant protein, 25
Regulation, 149
RESA , 67
Ribosomal RNA transcription, 111
mRNA, 281
5S rRNA gene, 295
5.8S rRNA gene, 111
RNA polymerase III, 295

Schistosoma mansoni, 199, 211
Signal peptide, 19
Specific binding, 221
Spectrin, 233
Sphingomyelin, 103

Tegument, 211
Tetrapeptide, 133
Theileria annulata, 253
Topoisomerase II, 139
Triacylglycerol, 175
Trichomonas vaginalis, 175
Tritrichomonas foetus, 175
Trypanosoma brucei, 139
Trypanosoma cruzi, 41, 75, 191, 245
Trypanosomatid, 75
Trypanosome, 139
Tubulin, 221